REPORTS ON THE FLORA OF THE BOSTON DISTRICT,—XXIX.

TEPHROSIA.

T. virginiana (L.) Pers. Dry sandy soil; well distributed but not very abundant.

TRIFOLIUM.

- T. AGRARIUM L. Fields and roadsides, common throughout.
- T. ARVENSE L. Dry sandy soil, very common throughout.
- T. DALMATICUM Vis. Dump, Dracut or Lowell (C. W. Swan, Aug. 6, 1884). Specimen in herb. N. E. Botanical Club. Adventive from Europe.
- T. Dubium Sibth. Waste land, S. Boston (C. E. Faxon, Sept. 30, 1878); sporadic in lawn, Hingham (C. H. Knowlton, June —, 1916).
- T. ECHINATUM Bieb. Wool-waste, Westford (Miss E. F. Fletcher, Rhodora xvii. 32, 1915). Native of southern Europe and Asia Minor.
 - T. HYBRIDUM L. Fields and roadsides, everywhere.
- T. Incarnatum L. Fields, occasionally persisting for one season after cultivation.
- T. Macrael Hook. & Arn. Parker River mills, Georgetown (Mrs. C. N. S. Horner, —, 1886); wool-waste, N. Chelmsford (W. P. Alcott, no date). Specimens in herb. N. E. Botanical Club. Native of western North America and Chili.
- T. MEDIUM L. Dry fields and pastures; many old specimens from Salem, Wenham and Danvers; also collected from Georgetown (E. F. Williams, July 10, 1901) and from Boston (Asa Gray, no date).
 - T. PRATENSE L. Fields and roadsides, everywhere.
 - T. PROCUMBENS L. Roadsides and fields, frequent.
- T. PURPUREUM Lois. Westford (Miss E. F. Fletcher, Rhodora xv. 172, 1913). Native of southern Europe and western Asia.
 - T. repens L. Fields and pastures everywhere.
 - T. RESUPINATUM L. South Boston (C. E. Perkins, Aug. 9, 1879).

Native of western Europe and the Mediterranean region. Specimen in herb. N. E. Botanical Club.

T. TOMENTOSUM L. Georgetown mills (Mrs. C. N. S. Horner, —, 1886). A very small specimen in herb. N. E. Botanical Club. Native of the Mediterranean region.

TRIGONELLA.

T. LACINIATA L. Charlestown (C. E. Perkins, July 4, 1879).

T. Noeana Boiss. (*T. Cassia* Boiss. of Dame & Collins, Flora Middlesex Co. 26, 1888). Waste-heap, Lowell (*C. W. Swan*, Aug. 4, 1884).

T. RAMOSA L. Charlestown (C. E. Perkins, July 4, 1879). The above specimens are in herb. N. E. Botanical Club.

VICIA.

V. ANGUSTIFOLIA Reichard. Waste ground, Lowell, Lynn, Welles-ley and S. Boston.

V. Angustifolia Reichard, var. segetalis (Thuill.) Koch. Waste places frequent. "On ye beach at Nehant" (M. Cutler, no date). Specimen in Gray Herb.

V. Angustifolia Reichard, var. uncinata (Desv.) R. & F. Dry gravel in field, Hingham (C. H. Knowlton, June 22, 1912).

V. Cracca L. Fields and roadsides, frequent.

V. Hirsuta (L.) S. F. Gray. Waste places, rare; Newburyport, W. Newbury, Danvers, Lowell, Dracut, Boston, Hingham, Wellesley.

V. SATIVA L. Waif in garden, Topsfield (Mrs. C. N. S. Horner, no date); Jamaica Plain (C. E. Faxon, —, 1882).

V. Tetrasperma (L.) Moench. Waste places and edges of woods, occasional.

V. VILLOSA Roth. Persistent after cultivation at Marshfield and Sherborn; perhaps elsewhere.

C. H. KNOWLTON | Committee on Walter Deane | Local Flora.